

# Towards an ECAL $e^-$ analysis paper?

## David Ward

- ❖ Would like to aim to progress from LCWS note to published paper(s). Envisage two papers:
  - ❖ 1. Hardware, calibration, technical performance (gain, noise, stability etc). Anne-Marie to coordinate.
  - ❖ 2. Electron response (linearity, resolution, uniformity etc.)
- ❖ Schedule – drafts by the end of 2007???
- ❖ Basic topics much as in the LCWS note.
- ❖ Several problems to be addressed before we can be ready. **Will mainly discuss these.**
- ❖ Comments on systematics.

# Content of analysis paper?


- ❖ Energy response
  - ❖ Linearity; uniformity (gap correction); angle dependence.
- ❖ Energy resolution
  - ❖ As a function of energy, angle, impact of gaps...
- ❖ Shower longitudinal profile
- ❖ Shower transverse profile
  - ❖ Effective Molière radius
- ❖ Position resolution
  - ❖ Using tracking
- ❖ Angular resolution
  - ❖ Using tracking
- ❖ 2-shower separation
  - ❖ Using double events/superimposed events
- ❖ All of the above compared with MC simulation.

# ECAL problem areas

- ❖ There are several problematic areas, where we see discrepancies between data and Monte Carlo.
- ❖ Some at least may be partly interconnected.
  - ❖ Low pulse height hits
  - ❖ Number of hits.
  - ❖ Interwafer gaps (effect of guard rings etc)
  - ❖ Transverse shower shape ( $\sim 10\%$  higher in data)
  - ❖ Shower depth (understanding of beam line, upstream material?)
  - ❖ Mismatch of energy scale ( $\sim 3\%$ ) between CERN and DESY


# Low energy hits

E Ecal hits /mips



- ❖ Pre3 processing agrees with MC quite well down to 1.2 MIPs
- ❖ Pre4 processing (SIPS correction) gives extra hits; agrees less well with MC.
- ❖ Effect grows with increasing energy.
- ❖ Anne-Marie's MC digitization and reconstruction doesn't have any significant effect (next slide...).
- ❖ Similar in 2007 data; possibly a bit worse


# Effect of MC digitization (45 GeV)



# Correlated with shower position


$N(<0.8\text{MIP})$

2006 Data 30 GeV



$N(<0.8\text{MIP})$


MC 30 GeV







# HitMap of hits below 0.8 MIP

HitMap ECAL Ehit<0.8MIP

2007 Data







# Number of hits (above threshold)













# Look at events in the high tail in data




Double shower.  
n.b. energy is normal for a single electron; if anything it's lower than average.



Such events are essentially not simulated at all in Mokka. There is much still to understand about the beam!


# Cut against double events

- ❖ Simple algorithm: form  $x$ - $y$  projection of all 30 layers.
- ❖ Apply Threshold  $T$
- ❖ Perform naive nearest-neighbour clustering of cells above threshold.
- ❖ Find  $T_{\max}$  below which two or more clusters first appear.
- ❖ Dramatic difference between data and MC
- ❖ Cut  $T_{\max} < 100$  to remove (most) double events.



# Compare with 2007 data (20 GeV)

Threshold for 2 clusters



2006


2007 e<sup>-</sup>  
Better, but  
other runs  
worse

2007 e<sup>+</sup>  
Much better,  
but still worse  
than MC

n.b. the e<sup>+</sup>  
beam is much  
narrower than  
the e<sup>-</sup> beam


# Effect of cut against double showers

N Ecal hits > Thresh




Helps with  $N_{\text{hits}}$ , but doesn't fully solve the problem.  
Little effect on  $E_{\text{tot}}$ .

Nhits




Etot




# Other energies?


N Ecal hits > Thresh




N Ecal hits > Thresh



N Ecal hits > Thresh




N Ecal hits > Thresh




Peak position vs. energy is still a issue

# Interwafer gaps; 30 GeV $e^-$

E vs  $\gamma$




E vs  $\gamma$



# Dip vs. Layer; 30 GeV $e^-$

Fit dip in energy vs  $y$  in each layer to a Gaussian.  
Plot fractional depth of dip, position and  $\sigma$ , for data and MC



Dip in MC still slightly deeper than in data


Beam inclined in data?  
Alignment issues?




# Radial energy distribution

Greatly improved by cut against double showers; at all CERN energies





E vs  $r_{hit}$



E vs  $r_{hit}$




# Shower depth



- Shower depth consistently » 0.2-0.3  $X_0$  deeper in MC than data.
- R.M.S. seems quite well modelled.
- Suggests problem with modelling the beam line rather than the material of ECAL?

# Shower depth – effect of double shower cut

Energy v X0







MC 7.78

Data 7.50

After T<sub>max</sub> cut 7.57


Not the main cause of the problem.

# Hits in Layer 1 only (30 GeV e<sup>-</sup>)




# Increase material in beam


Energy v X0




N Ecal hits layer 1



E Ecal /MIPs layer 1




E Ecal hits /MIPs layer 1




- Try increasing material in beam by 10% or 20%  $X_0$
- Technically done by doubling or trebling thickness of scintillators.
- Obviously too crude, but suggests ~ 10-20%  $X_0$  upstream material would be needed.
- Seems improbably much?
- Emphasises the need to understand the beam better.

# Uniformity across detector in 2007 data




# Another slice ( $40 < y < 50$ )




# Reweight two 20 GeV runs to same beam profile


Xav {Tmax<100}




Yav {Tmax<100}



Etot {Tmax<100}



Nhits {Tmax<100}




Take two 20 GeV runs (e<sup>-</sup> and e<sup>+</sup>) with similar beam position, but different width, and reweight to ~same profile.

Seems to be clear  
~1.5% difference in reconstructed energy.



# Energy scale CERN *c.f.* DESY

Glitch in data after correction for missing slabs; not simulated.



# Thoughts on systematic errors

- ❖ A major deficiency in what we've presented so far.
- ❖ Think what we might need to do, and whether software framework is adequate at present. Comments welcome.
  - ❖ Vary threshold cut.
    - ❖ Is the signal/noise cut in data reconstruction adequate?
  - ❖ Vary calibration constants within their statistical errors.
    - ❖ Reprocess using database? Or just smear? Or simulate using MC?
  - ❖ Vary cuts for selecting good  $e^\pm$  events.
  - ❖ Compare runs at same nominal energy.
  - ❖ Vary fit procedure/range for extracting response/resolution
  - ❖ Different inter-wafer gap corrections?
  - ❖ Intrinsic beam energy uncertainty ( $0.5\% \oplus 150 \text{ MeV}/p$ ) and spread (typically 0.5%).
- ❖ I think most of these could be done (or approximated well enough) using the existing reconstructed files.
- ❖ ...

# ...Continued...

- ❖ Would be good to add to the reconstructed files info about the beam (energy (nominal or “true”?), spread, position, angle, magnet currents?).
- ❖ Needs for the Monte Carlo?
  - ❖ Correct calorimeter geometry.
    - ❖ Including misalignments?
  - ❖ Dead cells (especially for 2007)
  - ❖ Signal-related crosstalk/pedestal shift, if we can understand the effect well enough.
  - ❖ More realistic simulation of the beam.
    - ❖ Spread of energy/position/angle; correlations between these.
    - ❖ Upstream material and showering.
    - ❖ Do we need a full beam line simulation?

# Summary

- ❖ Do we go for publication of 2006 data soon, or wait till we understand 2007 data?
- ❖ Should we split up topics into smaller papers?
- ❖ Do we combine the DESY and CERN data?
- ❖ Big questions to resolve:
  - ❖ Understand/simulate the beam more correctly
  - ❖ Treatment of energy loss in inter-wafer gaps, in a way that is angle-independent.
  - ❖ Characterise/correct/simulate “square events” and other coherent signal-related effects, e.g. SIPS.
- ❖ Think realistically about systematics. We are not making precision SM measurements, but we do have to assess the reliability of anything we measure.