Antiprotons at Fermilab: New Directions in Hyperon, Charm, and Antimatter Physics

Daniel M. Kaplan

ILLINOIS INSTITUTE OF TECHNOLOGY Transforming Lives. Inventing the Future. www.iit.edu



High Energy Physics Seminar Imperial College London 24 November 2008

Outline

(Varied menu!)

- Hyperon CP violation
- Low-energy antiprotons
- A new experiment
- Issues in charmonium
- Charm mixing
- Antihydrogen measurements
- Summary

• An old topic:

• An old topic:

PHYSICAL REVIEW

VOLUME 184, NUMBER 5

25 AUGUST 1969

Final-State Interactions in Nonleptonic Hyperon Decay

O. E. OVERSETH*

The University of Michigan, Ann Arbor, Michigan 48104

AND

S. PAKVASA[†] University of Hawaii, Honolulu, Hawaii 96822 (Received 1 April 1969)

E. Tests for CP and CPT Invariance

Thus in hyperon decay, $\bar{\alpha} \neq -\alpha$ implies *CP* violation in this process independent of the validity of the *CPT* theorem. This is also true if $\bar{\beta} \neq -\beta$.

Also, as usual, CPT invariance implies equality of Λ^0 and $\overline{\Lambda}^0$ lifetimes, whereas CP invariance implies equality of partial rates $\Gamma^0 = \overline{\Gamma}^0$, and $\Gamma^- = \overline{\Gamma}^+$. This is also true when final-state interactions are included in the analysis.





• CP-odd observables:

- CP-odd observables:
 - ► Decay amplitude for $\Delta S = 1$ decay of spin-1/2 strange baryon into spin-1/2 baryon and meson (e.g., $\Lambda \rightarrow p \pi^-$):

$$M = S + P\vec{\sigma} \cdot \hat{q}_p$$

- CP-odd observables:
 - ► Decay amplitude for $\Delta S = 1$ decay of spin-1/2 strange baryon into spin-1/2 baryon and meson (e.g., $\Lambda \rightarrow p \pi^-$):

$$M = S + P\vec{\sigma} \cdot \hat{q}_p$$

S-wave amplitude

- CP-odd observables:
 - ► Decay amplitude for $\Delta S = 1$ decay of spin-1/2 strange baryon into spin-1/2 baryon and meson (e.g., $\Lambda \rightarrow p \pi^-$):



- CP-odd observables:
 - ► Decay amplitude for $\Delta S = 1$ decay of spin-1/2 strange baryon into spin-1/2 baryon and meson (e.g., $\Lambda \rightarrow p \pi^-$):



- CP-odd observables:
 - ► Decay amplitude for $\Delta S = 1$ decay of spin-1/2 strange baryon into spin-1/2 baryon and meson (e.g., $\Lambda \rightarrow p \pi^-$):



- CP-odd observables:
 - ► Decay amplitude for $\Delta S = 1$ decay of spin-1/2 strange baryon into spin-1/2 baryon and meson (e.g., $\Lambda \rightarrow p \pi^-$):



- CP-odd observables:
 - ► Decay amplitude for $\Delta S = 1$ decay of spin-1/2 strange baryon into spin-1/2 baryon and meson (e.g., $\Lambda \rightarrow p \pi^-$):



- CP-odd observables:
 - ► Decay amplitude for $\Delta S = 1$ decay of spin-1/2 strange baryon into spin-1/2 baryon and meson (e.g., $\Lambda \rightarrow p \pi^-$):



$$\Rightarrow A_{\Lambda} = \frac{\alpha_{\Lambda} + \overline{\alpha}_{\Lambda}}{\alpha_{\Lambda} - \overline{\alpha}_{\Lambda}}, B_{\Lambda} = \frac{\beta_{\Lambda} + \overline{\beta}_{\Lambda}}{\beta_{\Lambda} - \overline{\beta}_{\Lambda}}, \Delta_{\Lambda} = \frac{\Gamma_{\Lambda \to P\pi} - \overline{\Gamma}_{\Lambda \to P\pi}}{\Gamma_{\Lambda \to P\pi} + \overline{\Gamma}_{\Lambda \to P\pi}} CP-odd$$

D.M. Kaplan, IIT New Experiments with Antiprotons ICL HEP Seminar

$$\Rightarrow A_{\Lambda} \equiv \frac{\alpha_{\Lambda} + \overline{\alpha}_{\Lambda}}{\alpha_{\Lambda} - \overline{\alpha}_{\Lambda}}, \ B_{\Lambda} \equiv \frac{\beta_{\Lambda} + \overline{\beta}_{\Lambda}}{\beta_{\Lambda} - \overline{\beta}_{\Lambda}}, \ \Delta_{\Lambda} \equiv \frac{\Gamma_{\Lambda \to P\pi} - \overline{\Gamma}_{\Lambda \to P\pi}}{\Gamma_{\Lambda \to P\pi} + \overline{\Gamma}_{\Lambda \to P\pi}} \text{ CP-odd}$$

Hyperon CP Violation?

$$\Rightarrow A_{\Lambda} = \frac{\alpha_{\Lambda} + \overline{\alpha}_{\Lambda}}{\alpha_{\Lambda} - \overline{\alpha}_{\Lambda}}, B_{\Lambda} = \frac{\beta_{\Lambda} + \overline{\beta}_{\Lambda}}{\beta_{\Lambda} - \overline{\beta}_{\Lambda}}, \Delta_{\Lambda} = \frac{\Gamma_{\Lambda \to P\pi} - \overline{\Gamma}_{\Lambda \to P\pi}}{\Gamma_{\Lambda \to P\pi} + \overline{\Gamma}_{\Lambda \to P\pi}} \text{ CP-odd}$$
• $p \neq distribution in \Lambda rest frame: \frac{dN}{d\cos\theta} = 1 + \alpha_{\Lambda}P_{\Lambda}\cos\theta$

$$= \int_{1}^{1} \int_{0}^{0} \int_{0}^{1} \int_{0$$

Hyperon CP Violation?

$$\Rightarrow A_{\Lambda} = \frac{\alpha_{\Lambda} + \overline{\alpha}_{\Lambda}}{\alpha_{\Lambda} - \overline{\alpha}_{\Lambda}}, B_{\Lambda} = \frac{\beta_{\Lambda} + \overline{\beta}_{\Lambda}}{\beta_{\Lambda} - \overline{\beta}_{\Lambda}}, \Delta_{\Lambda} = \frac{\Gamma_{\Lambda \to P\pi} - \overline{\Gamma}_{\Lambda \to P\pi}}{\Gamma_{\Lambda \to P\pi} + \overline{\Gamma}_{\Lambda \to P\pi}} \text{ CP-odd}$$
• $p \ \ distribution in \ \Lambda rest frame: \frac{dN}{d \cos \theta} = 1 + \alpha_{\Lambda} P_{\Lambda} \cos \theta$
• $p \ \ distribution in \ \Lambda rest frame: \frac{dN}{d \cos \theta} = 1 + \alpha_{\Lambda} P_{\Lambda} \cos \theta$
• $CP \ conserved \Leftrightarrow slope = -\overline{slope}$

• For precise measurement of A, need excellent knowledge of relative Λ and $\overline{\Lambda}$ polarizations!

- For precise measurement of A, need excellent knowledge of relative Λ and $\overline{\Lambda}$ polarizations!
 - →HyperCP "trick": $\Xi^- \rightarrow \Lambda \pi^-$ decay gives $P_{\Lambda} = -P_{\overline{\Lambda}}$



New Experiments with Antiprotons

- For precise measurement of A, need excellent knowledge of relative Λ and $\overline{\Lambda}$ polarizations!
 - →HyperCP "trick": $\Xi^- \rightarrow \Lambda \pi^-$ decay gives $P_{\Lambda} = -P_{\overline{\Lambda}}$



• Unequal slopes \Rightarrow CP violated!

• Theory & experiment:

$$\alpha_{\overline{\Lambda}} = -\alpha_{\Lambda}$$

• One can form a CP violating asymmetry

$$A_{\Lambda} = \frac{\alpha_{\Lambda} + \alpha_{\overline{\Lambda}}}{\alpha_{\Lambda} + \alpha_{\overline{\Lambda}}}$$

- Theory & experiment:
- Search for direct CP violation in Λ decay ..., e.g., PRL 55, 162 (1985); PRD 34, 833 Need to produce Λ , $\overline{\Lambda}$ with known pola $|A_{\Xi\Lambda}| < 5 \times 10^{-5}$ (1986); PLB 272, 411 (1991)] Experiment Decay Mode A_{Λ} 056001 (2003)]
 - **R608 at ISR** $pp \to \Lambda X, \bar{p}p \to \bar{\Lambda}X$ -0.02 \pm 0.14
 - DM2 at Orsay $e^+e^- \rightarrow J/\Psi \rightarrow \Lambda \bar{\Lambda}$ 0.01 \pm 0.10
 - **PS185 at LEAR** $p\bar{p} \to \Lambda\bar{\Lambda}$ **PS185 at LEAR** [P. Chauvat et al., PL 163B (1985) 273] **-0.013 ± 0.022**73

[M.H. Tixier et al., PL B212 (1988) 523] 273

tion?

E756 at Fermilab	$\Xi ightarrow \Lambda \pi, \Lambda ightarrow p\pi$	0.012 ± 0.014 [K.B. Luk et al., PRL 85, 4860 (2000)]
	$\Xi \to \Lambda \pi, \Lambda \to p \pi$	
E756 at Fermilab	$\Xi ightarrow \Lambda \pi, \Lambda ightarrow p\pi$	0.012 ± 0.014

ICL HEP Seminar 8

$$\alpha_{\overline{\Lambda}} = -\alpha_{\Lambda}$$

• One can form a CP violating asymmetry

$$A_{\Lambda} = \frac{\alpha_{\Lambda} + \alpha_{\overline{\Lambda}}}{\alpha_{\Lambda} = \alpha_{\overline{\Lambda}}}$$

• Theory $\hat{\&} \exp{\hat{e}rim\hat{e}nt}$:

Search for direct	t CP violation in Λ de	$(ay)^{-1}$, e.g., PRL 55, 162 (1985); PRD 34, 833	
Need to produce Λ , $\bar{\Lambda}$ with known pola $ A_{F\Lambda} < 5 \times 10^{-5}$ [J. Tandean, G. Valencia, Phys. Rev. D 67,			
Experiment	Decay Mode	$A_{\Lambda} = 056001 (2003)]$	
R608 at ISR Experiment	$pp \to \Lambda X, \bar{p}p \to \bar{\Lambda}X$ Decay Mode	$\begin{array}{c} \textbf{-0.02} \pm \textbf{0.14} \\ \textbf{A}_{\Lambda} \end{array}$	
R608 at ISR	$pp \to \Lambda X, \bar{p}p \to \bar{\Lambda} X$	-0.02 ± 0.14 [P. Chauyat et al.; BL 163B (1985) 273]	
DM2 at Orsay	$e^+e^- \to J/\Psi \to \Lambda \bar{\Lambda}$	0.01 ± 0.10 [M.H. Tixier et al.; PL B212 (1988) 523]	
PS185 at LEAR	$p\bar{p} \rightarrow \Lambda \bar{\Lambda}$	0.006 ± 0.015 [P.D. Barnes et al.; NP B 36A (1997) 46]	
Sequence Experiment	Decay Mode	$\mathbf{A}_{\Xi} + \mathbf{A}_{\Lambda}$	
E756 at Fermilab	$\Xi \rightarrow \Lambda \pi, \Lambda \rightarrow p\pi$	0.012 ± 0.014 [K.B. Luk et al., PRL 85, 4860 (2008)]	
E871 at Fermilab	Ē⇒₳₳,₳⇒₽₹	$(0.0 \pm 6.7) \times 10^{-4}$ [T. Holmstrom et al., PRL 93. 262001 (2004)]	
(HyperCP) E7 56 at Ferm ilab	$\Xi ightarrow \Lambda \pi, \Lambda ightarrow p\pi$	$(6 \pm 2 \pm 2) \times 10^{-4}$ [BEACH08 preliminary]	
		ICL HEP Seminar	

tion?





Made possible by...









...and Fast HyperCP DAQ System

...and Fast HyperCP DAQ System

\approx 20,000 channels of MWPC latches


...and Fast HyperCP DAQ System

\approx 20,000 channels of MWPC latches



\approx 100 kHz of triggers

...and Fast HyperCP DAQ System

\approx 20,000 channels of MWPC latches



 \approx 100 kHz of triggers ...written to 32 tapes in parallel



New Experiments with Antiprotons

ICL HEP Seminar II

HyperCP Collaboration



A. Chan, Y.-C. Chen, C. Ho, P.-K. Teng Academia Sinica, Taiwan

K. Clark, M. Jenkins University of South Alabama, USA

W.-S. Choong, Y. Fu, G. Gidal, T. D. Jones, K.-B. Luk*, P. Gu, P. Zyla University of California, Berkeley, USA

> C. James, J. Volk Fermilab, USA

J. Felix, G. Moreno, M. Sosa University of Guanajuato, Mexico

R. Burnstein, A. Chakravorty, D. Kaplan, L. Lederman, D. Rajaram, H. Rubin, N. Solomey, C. White *Illinois Institute of Technology, USA*

N. Leros, J.-P. Perroud University of Lausanne, Switzerland

H. R. Gustafson, M. Longo, F. Lopez, H. Park University of Michigan, USA

E. C. Dukes*, C. Durandet, T. Holmstrom, M. Huang, L. C. Lu, K. S. Nelson University of Virginia, USA *co-sp

*co-spokespersons

D. M. Kaplan, IIT

New Experiments with Antiprotons

ICL HEP Seminar 12

- $\phi_{\Xi} = (-2.39 \pm 0.64 \pm 0.64)^{\circ} \Rightarrow \beta_{\Xi} \neq 0$ 2nd non-zero transverse asymm.
- $\alpha_{\Omega} = 0.0175 \pm 0.0024 \neq 0 \qquad \Rightarrow \Omega^{-} \rightarrow \Lambda K^{-}$ violates parity

- $\phi_{\Xi} = (-2.39 \pm 0.64 \pm 0.64)^{\circ} \Rightarrow \beta_{\Xi} \neq 0$ 2nd non-zero transverse asymm.
- $\alpha_{\Omega} = 0.0175 \pm 0.0024 \neq 0 \qquad \Rightarrow \Omega^{-} \rightarrow \Lambda K^{-}$ violates parity
- $\frac{1}{2}[\alpha(\Lambda K^{-}) + \alpha(\overline{\Lambda} K^{+})] = -0.004 \pm 0.040$ (but conserves CP)

- $\phi_{\Xi} = (-2.39 \pm 0.64 \pm 0.64)^{\circ} \Rightarrow \beta_{\Xi} \neq 0$ 2nd non-zero transverse asymm.
- $\alpha_{\Omega} = 0.0175 \pm 0.0024 \neq 0 \qquad \Rightarrow \Omega^{-} \rightarrow \Lambda K^{-}$ violates parity
- $\frac{1}{2}[\alpha(\Lambda K^{-}) + \alpha(\overline{\Lambda} K^{+})] = -0.004 \pm 0.040$ (but conserves CP)
- $A_{\Xi\Lambda} = (0.0 \pm 5.1 \pm 4.4) \times 10^{-4} \implies \Xi^- \rightarrow \Lambda \pi^- \text{ conserves CP}$

- $\phi_{\Xi} = (-2.39 \pm 0.64 \pm 0.64)^{\circ} \Rightarrow \beta_{\Xi} \neq 0$ 2nd non-zero transverse asymm.
- $\alpha_{\Omega} = 0.0175 \pm 0.0024 \neq 0 \qquad \Rightarrow \Omega^{-} \rightarrow \Lambda K^{-}$ violates parity
- $\frac{1}{2}[\alpha(\Lambda K^{-}) + \alpha(\overline{\Lambda} K^{+})] = -0.004 \pm 0.040$ (but conserves CP)
- $A_{\Xi\Lambda} = (0.0 \pm 5.1 \pm 4.4) \times 10^{-4} \Rightarrow \Xi^- \rightarrow \Lambda \pi^- \text{ conserves CP}$ (1st $\approx 5\%$ of sample - full analysis still in progress)

- $\phi_{\Xi} = (-2.39 \pm 0.64 \pm 0.64)^{\circ} \Rightarrow \beta_{\Xi} \neq 0$ 2nd non-zero transverse asymm.
- $\alpha_{\Omega} = 0.0175 \pm 0.0024 \neq 0 \qquad \Rightarrow \Omega^{-} \rightarrow \Lambda K^{-}$ violates parity
- $\frac{1}{2}[\alpha(\Lambda K^{-}) + \alpha(\overline{\Lambda} K^{+})] = -0.004 \pm 0.040$ (but conserves CP)
- $A_{\Xi\Lambda} = (0.0 \pm 5.1 \pm 4.4) \times 10^{-4} \Rightarrow \Xi^- \rightarrow \Lambda \pi^- \text{ conserves CP}$ (1st $\approx 5\%$ of sample - full analysis still in progress)
- $\Sigma^+ \rightarrow p \mu^+ \mu_{\overline{\cdot}}$ smallest baryon BR ever seen!

- $\phi_{\Xi} = (-2.39 \pm 0.64 \pm 0.64)^{\circ} \Rightarrow \beta_{\Xi} \neq 0$ 2nd non-zero transverse asymm.
- $\alpha_{\Omega} = 0.0175 \pm 0.0024 \neq 0 \qquad \Rightarrow \Omega^{-} \rightarrow \Lambda K^{-}$ violates parity
- $\frac{1}{2}[\alpha(\Lambda K^{-}) + \alpha(\overline{\Lambda} K^{+})] = -0.004 \pm 0.040$ (but conserves CP)
- $A_{\Xi\Lambda} = (0.0 \pm 5.1 \pm 4.4) \times 10^{-4} \Rightarrow \Xi^- \rightarrow \Lambda \pi^- \text{ conserves CP}$ (1st $\approx 5\%$ of sample - full analysis still in progress)
- $\Sigma^+ \rightarrow p \mu^+ \mu^- : \mathcal{B} \approx 9 \times 10^{-8} \text{ (or } 3 \times 10^{-8} \text{ if intermediate } P^0)$







PRL 98, 081802 (2007)

Does the HyperCP Evidence for the Decay $\Sigma^+ \rightarrow p \mu^+ \mu^-$ Indicate a Light Pseudoscalar Higgs Boson?

Xiao-Gang He*

Department of Physics and Center for Theoretical Sciences, National Taiwan University, Taipei, Taiwan

Jusak Tandean[†]

Departments of Mathematics, Physics, and Computer Science, University of La Verne, La Verne, California 91750, USA

G. Valencia[‡]

Department of Physics and Astronomy, Iowa State University, Ames, Iowa 50011, USA (Received 2 November 2006; published 22 February 2007)

The HyperCP Collaboration has observed three events for the decay $\Sigma^+ \rightarrow p\mu^+\mu^-$ which may be interpreted as a new particle of mass 214.3 MeV. However, existing data from kaon and *B*-meson decays provide stringent constraints on the construction of models that support this interpretation. In this Letter we show that the "HyperCP particle" can be identified with the light pseudoscalar Higgs boson in the next-to-minimal supersymmetric standard model, the A_1^0 . In this model there are regions of parameter space where the A_1^0 can satisfy all the existing constraints from kaon and *B*-meson decays and mediate $\Sigma^+ \rightarrow p\mu^+\mu^-$ at a level consistent with the HyperCP observation. PRL 98, 081802 (2007)

Does the HyperCP Evidence for the Decay $\Sigma^+ \rightarrow p \mu^+ \mu^-$ Indicate a Light Pseudoscalar Higgs Boson?

Xiao-Gang He*

Department of Physics and Center for Theoretical Sciences, National Taiwan University, Taipei, Taiwan

Jusak Tandean[†]

Departments of Mathematics, Physics, and Computer Science, University of La Verne, La Verne, California 91750, USA

G. Valencia[‡]

Department of Physics and Astronomy, Iowa State University, Ames, Iowa 50011, USA (Received 2 November 2006; published 22 February 2007)

The HyperCP Collaboration has observed three events for the decay $\Sigma^+ \rightarrow p\mu^+\mu^-$ which may be interpreted as a new particle of mass 214.3 MeV. However, existing data from kaon and *B*-meson decays provide stringent constraints on the construction of models that support this interpretation. In this Letter we show that the "HyperCP particle" can be identified with the light pseudoscalar Higgs boson in the next-to-minimal supersymmetric standard model, the A_1^0 . In this model there are regions of parameter space where the A_1^0 can satisfy all the existing constraints from kaon and *B*-meson decays and mediate $\Sigma^+ \rightarrow p\mu^+\mu^-$ at a level consistent with the HyperCP observation.

- Holmstrom et al., PRL **93**, 26201 (2004):
 - analysis of \approx 5% of Ξ^- sample, 10% of Ξ^+

Ξ[±] CPViolation

- Holmstrom et al., PRL **93**, 26201 (2004):
 - analysis of \approx 5% of Ξ^- sample, 10% of Ξ^+



• Holmstrom et al., PRL **93**, 26201 (2004):

- analysis of \approx 5% of Ξ^- sample, 10% of Ξ^+



After weighting events to correct for unequal production spectra, etc.: $\delta(\cos\theta \text{ slope}) = 0$

• Holmstrom et al., PRL **93**, 26201 (2004):

- analysis of \approx 5% of Ξ^- sample, 10% of Ξ^+



After weighting events to correct for unequal production spectra, etc.: $\delta(\cos\theta \text{ slope}) = 0$

• C. Materniak, BEACH08:

• Holmstrom et al., PRL **93**, 26201 (2004):

- analysis of \approx 5% of Ξ^- sample, 10% of Ξ^+



• Holmstrom et al., PRL **93**, 26201 (2004):

- analysis of \approx 5% of Ξ^- sample, 10% of Ξ^+



- Tevatron fixed-target is no more
- CERN fixed-target not as good (energy, duty factor)
- Main Injector fixed-target not as good (same reasons)
- AND HyperCP was already rate-limited

- Tevatron fixed-target is no more
- CERN fixed-target not as good (energy, duty factor)
- Main Injector fixed-target not as good (same reasons)
- AND HyperCP was already rate-limited



- Tevatron fixed-target is no more
- CERN fixed-target not as good (energy, duty factor)
- Main Injector fixed-target not as good (same reasons)
- AND HyperCP was already rate-limited
- Big collider experiments can't trigger efficiently



- Tevatron fixed-target is no more
- CERN fixed-target not as good (energy, duty factor)
- Main Injector fixed-target not as good (same reasons)
- AND HyperCP was already rate-limited
- Big collider experiments can't trigger efficiently





• Until "HyperCP era," world's best limit on hyperon CP violation came from PS185 at LEAR:

 Until "HyperCP era," world's best limit on hyperon CP violation came from PS185 at LEAR:

Experiment	Decay Mode	\mathbf{A}_{Λ}
R608 at ISR	$pp \to \Lambda X, \bar{p}p \to \bar{\Lambda} X$	-0.02 ± 0.14 [P. Chauvat et al., PL 163B (1985) 273
DM2 at Orsay	$e^+e^- \to J/\Psi \to \Lambda \bar{\Lambda}$	0.01 ± 0.10 [M.H. Tixier et al., PL B212 (1988) 523
PS185 at LEAR	$p \bar{p} ightarrow \Lambda ar{\Lambda}$	0.006 ± 0.015 [P.D. Barnes et al., NP B 56A (1997) 4
Experiment	Decay Mode	$\mathbf{A}_{\Xi} + \mathbf{A}_{\Lambda}$
E756 at Fermilab	$\Xi ightarrow \Lambda \pi, \Lambda ightarrow p\pi$	0.012 ± 0.014 [K.B. Luk et al., PRL 85, 4860 (2000)]
E871 at Fermilab (HyperCP)	$\Xi \to \Lambda \pi, \Lambda \to p\pi$	$(0.0 \pm 6.7) \times 10^{-4}$ [T. Holmstrom et al., PRL 93. 262001 (2004)]
		$(6 \pm 2 \pm 2) \times 10^{-4}$ [BEACH08 preliminary]

 Until "HyperCP era," world's best limit on hyperon CP violation came from PS185 at LEAR:

Experiment	Decay Mode	\mathbf{A}_{Λ}	
R608 at ISR	$pp \to \Lambda X, \bar{p}p \to \bar{\Lambda} X$	-0.02 ± 0.14 [P. Cl	hauvat et al., PL 163B (1985) 273]
DM2 at Orsay	$e^+e^- \to J/\Psi \to \Lambda \bar{\Lambda}$	0.01 ± 0.10 [M.H	. Tixier et al., PL B212 (1988) 523]
PS185 at LEAR	$par{p} o \Lambdaar{\Lambda}$	0.006 ± 0.015 [P.D.	Barnes et al., NP B 56A (1997) 46
Experiment	Decay Mode	$\mathbf{A}_{\Xi} + \mathbf{A}_{\Lambda}$	
E756 at Fermilab	$\Xi ightarrow \Lambda \pi, \Lambda ightarrow p\pi$	0.012 ± 0.014 [K.B.	Luk et al., PRL 85, 4860 (2000)]
E871 at Fermilal (HyperCP)	$\Xi \to \Lambda \pi, \Lambda \to p\pi$	$(0.0 \pm 6.7) \times 10^{-4}$ [7]	^c . Holmstrom et al., RL 93. 262001 (2004)]
		$(6 \pm 2 \pm 2) \times 10^{-4}$ [B	BEACH08 preliminary]
- UT		····	

• PS185 was limited by LEAR \overline{p} flux ($\leq 10^{5}/s$)

• PS185 was limited by LEAR \overline{p} flux ($\leq 10^{5}/s$)



18

• PS185 was limited by LEAR \overline{p} flux ($\leq 10^{5}/s$)



18

- Also good for charmonium:
 - Thanks to superb precision of antiproton beam energy and momentum spread, E760/835 @ Fermilab Antiproton Accumulator made very precise measurements of charmonium parameters, e.g.:
• Also good for charmonium:

Thanks to superb precision of antiproton beam energy and momentum spread, E760/835 @ Fermilab Antiproton Accumulator made very precise

measurements of charmonium

parameters, e.g.:

 $\chi_{c0}(1P)$ MASS

VALUE (MeV) EVTS		DOCUMENT ID		TECN	COMMENT		
3414.76±	0.35	OUR A	/ERAGE	Error includes sca	ale fa	ctor of	1.2.
$3414.21\pm$	0.39	± 0.27		ABLIKIM	05 G	BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$
3414.7 $\stackrel{+}{_}$	0.7 0.6	± 0.2		¹ ANDREOTTI	03	E835	$\overline{p}p \rightarrow \chi_{c0} \rightarrow \pi^0 \pi^0$
3415.5 \pm	0.4	± 0.4	392	² BAGNASCO	02	E835	$\overline{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
3417.4 +	1.8 1.9	± 0.2		¹ AMBROGIANI	99 B	E835	$\overline{p}p \rightarrow e^+e^-\gamma$
$\begin{array}{rrrr} 3414.1 & \pm \\ 3417.8 & \pm \end{array}$	0.6 0.4	±0.8 ±4		BAI ¹ GAISER	99в 86	BES CBAL	$\psi(2S) ightarrow \gamma X \ \psi(2S) ightarrow \gamma X$

 $I^{G}(J^{PC}) = 0^{+}(0^{++})$

New Experiments with Antiprotons

• Also good for charmonium:

Thanks to superb precision of antiproton beam energy and momentum spread, E760/835 @ Fermilab Antiproton Accumulator made very precise

measurements of charmonium parameters, e.g.:

$\chi_{c0}(1P)$
$\chi_{c2}(1P)$

$$G(J^{PC}) = 0^+(0^{++})$$



$$I^{G}(J^{PC}) = 0^{+}(2^{+})$$

See the Review on " $\psi(2S)$ and χ_c branching ratios" before the $\chi_{c0}(1P)$ Listings.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3556.20 ± 0.09 OUR	AVERAGE			
$3555.70~\pm~0.59~\pm0.39$	9	ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma \chi_{c2}$
$3556.173 \pm 0.123 \pm 0.02$	20	ANDREOTTI	05A E835	$p \overline{p} \rightarrow e^+ e^- \gamma$
3559.9 ± 2.9		EISENSTEIN	01 CLE2	$e^+e^{\perp} \rightarrow$
				$e^+e^-\chi_{c2}$
3556.4 ± 0.7		BAI	99b BES	$\psi(2S) ightarrow \gamma X$
$3556.22 \pm 0.131 \pm 0.02$ New Experiments with A	20 585 Intiprotons	¹ ARMSTRONG	92 E760 ICL F	$\begin{array}{c c} \overline{p}p \rightarrow e^+ e^- \gamma \\ \hline \text{IEP Seminar} & 19 \end{array}$

D. M. Kaplan, IIT

- Also good for charmonium:
 - Thanks to superb precision of antiproton beam energy and momentum spread, E760/835 @ Fermilab Antiproton Accumulator made very precise measurements of charmonium parameters, e.g.:
 - best measurements of various η_c, χ_c, h_c masses, widths, branching ratios,...
 - interference of continuum & resonance signals
- GSI-Darmstadt upgrading to similar facility, done ≈2015
 D. M. Kaplan, IIT New Experiments with Antiprotons ICL HEP Seminar 20

Fermilab Antiproton Source is world's highest-energy and most intense

	Stack	Clock Hours	$\overline{\mathbf{p}}/\mathbf{Yr}$	
	Rate $(10^{10}/hr)$	Duty Factor	/ Yr	(10^{13})
CERN AD			3800	0.4
FNAL (Accumulator)	20	15%	5550	17
FNAL (New Ring)	20	90%	5550	100
FAIR (≥2015)	3.5	90%	2780	9

Table I: Antiproton Intensities at Existing and Future Facilities

...even after FAIR turns on

- Once Tevatron shuts down (\approx 2010),
 - Reinstall E835 EM spectrometer

- Once Tevatron shuts down (\approx 2010),
 - Reinstall E835 EM spectrometer



New Experiments with Antiprotons

- Once Tevatron shuts down (≈ 2010),
 - Reinstall E835 EM spectrometer
 - Add small magnetic spectrometer



- Once Tevatron shuts down (\approx 2010),
 - Reinstall E835 EM spectrometer
 - Add small magnetic spectrometer



- Once Tevatron shuts down (\approx 2010),
 - Reinstall E835 EM spectrometer
 - Add small magnetic spectrometer
 - Add TOF system



- Once Tevatron shuts down (\approx 2010),
 - Reinstall E835 EM spectrometer
 - Add small magnetic spectrometer
 - Add TOF system
 - Add wire or pellet target



- Once Tevatron shuts down (\approx 2010),
 - Reinstall E835 EM spectrometer
 - Add small magnetic spectrometer
 - Add TOF system
 - Add wire or pellet target
 - Add 2^{ndary}-vertex trigger



Return Yoke

pBEAM

INNER DETECTOR

<\$10M

Return Yoke

()

SciFi

LUMINOSITY

ICL HEP Seminar 22

SiPix

- Once Tevatron shuts down (\approx 2010),
 - Reinstall E835 EM spectrometer
 - Add small magnetic spectrometer
 - Add TOF system
 - Add wire or pellet target
 - Add 2^{ndary}-vertex trigger

pBEAM

<\$10M

Return Yoke

2000000

Scil

LUMINOSITY

ICL HEP Seminar 22

SiPix

- Once Tevatron shuts down (\approx 2010),
 - Reinstall E835 EM spectrometer
 - Add small magnetic spectrometer
 - Add TOF system
 - Add wire or pellet target
 - Add 2^{ndary}-vertex trigger
 - Run $p_{\overline{p}} = 5.4 \text{ GeV}/c \ (2m_{\Omega} < \sqrt{s} < 2m_{\Omega} + m_{\pi_0})$ @ $\mathcal{L} \sim 10^{32} \text{ cm}^{-2} \text{ s}^{-1} \ (10 \times \text{E835})$

pBEAM

<\$10M

Return Yoke

()

SciFi

LUMINOSITY

ICL HEP Seminar 22

SiPix

- Once Tevatron shuts down (\approx 2010),
 - Reinstall E835 EM spectrometer
 - Add small magnetic spectrometer
 - Add TOF system
 - Add wire or pellet target
 - Add 2^{ndary}-vertex trigger
 - Run $p_{\overline{p}} = 5.4 \text{ GeV}/c (2m_{\Omega} < \sqrt{s} < 2m_{\Omega} + m_{\pi_0})$ @ $\mathcal{L} \sim 10^{32} \text{ cm}^{-2} \text{ s}^{-1} (10 \times \text{E835})$

 $\Rightarrow \sim 10^8 \Omega^- \overline{\Omega}^+/yr$

pBEAM

<\$10M

Return Yoke

()

2000000

ICL HEP Seminar 22

SciFi

SiPix

- Once Tevatron shuts down (\approx 2010),
 - Reinstall E835 EM spectrometer
 - Add small magnetic spectrometer
 - Add TOF system
 - Add wire or pellet target
 - Add 2^{ndary}-vertex trigger
 - Run $p_{\overline{p}} = 5.4 \text{ GeV/c} (2m_{\Omega} < \sqrt{s} < 2m_{\Omega} + m_{\pi_0})$ @ $\mathcal{L} \sim 10^{32} \text{ cm}^{-2} \text{ s}^{-1} (10 \times \text{E835})$

 $\Rightarrow \sim 10^8 \Omega^- \overline{\Omega}^+/yr + \sim 10^{12}$ inclusive hyperon events!

• Observe many more $\Sigma^+ \to p \mu^+ \mu^-$ events and confirm or refute SUSY interpretation

- Observe many more $\Sigma^+ \rightarrow p \mu^+ \mu^-$ events and confirm or refute SUSY interpretation
- Discover or limit $\Omega^- \rightarrow \Xi^- \mu^+ \mu^-$ and confirm or refute SUSY interpretation

- Observe many more $\Sigma^+ \rightarrow p \mu^+ \mu^-$ events and confirm or refute SUSY interpretation
- Discover or limit $\Omega^- \rightarrow \Xi^- \mu^+ \mu^-$ and confirm or refute SUSY interpretation Predicted $\mathcal{B} \sim 10^{-6}$

if P^0 real

- Observe many more $\Sigma^+ \rightarrow p \mu^+ \mu^-$ events and confirm or refute SUSY interpretation
- Discover or limit $\Omega^- \rightarrow \Xi^- \mu^+ \mu^-$ and confirm or refute SUSY interpretation Predicted $\mathcal{B} \sim 10^{-6}$
- Discover or limit *CP* violation in $\Omega^- \to \Lambda K^$ and $\Omega^- \to \Xi^0 \pi^-$ via partial-rate asymmetries

if P^0 real

- Observe many more $\Sigma^+ \rightarrow p \mu^+ \mu^-$ events and confirm or refute SUSY interpretation
- Discover or limit $\Omega^- \rightarrow \Xi^- \mu^+ \mu^-$ and confirm or refute SUSY interpretation Predicted $\mathcal{B} \sim 10^{-6}$
- Discover or limit *CP* violation in $\Omega^- \to \Lambda K^$ and $\Omega^- \to \Xi^0 \pi^-$ via partial-rate asymmetries

Predicted $\Delta \mathcal{B} \sim 10^{-5}$ in SM, $\leq 10^{-3}$ if NP if P^0 real

 Much interest lately in new states observed in charmonium region: X(3872), X(3940), Y(3940), Y(4260), and Z(3930)

- Much interest lately in new states observed in charmonium region: X(3872), X(3940), Y(3940), Y(4260), and Z(3930)
- X(3872) of particular interest b/c may be the first hadron-antihadron ($D^0 \overline{D}^{*0}$ + c.c.) molecule

• Belle, Aug. 2003: $B^{\pm} \longrightarrow X + K^{\pm}, X \longrightarrow J/\psi \pi^{+}\pi^{-}$



- Since confirmed by CDF, D0, & BaBar
- Not consistent with being charmonium state
- Very near $D^0 \overline{D}^{*0}$ threshold ($\Delta mc^2 = -0.35 \pm 0.69$ MeV)

XYZ hadronic transitions

• Many new states : ?

State	EXP	М + і Г (MeV)	J ^{PC}	Decay Modes Observed	Production Modes Observed
X(3872)	Belle,CDF, DO, Cleo, BaBar	3871.2±0.5 + i(<2.3)	1++	π⁺π⁻J/ψ, π⁺π⁻π⁰J/ψ, ƳJ/ψ	B decays, ppbar
	Belle BaBar	3875.4±0.7 ^{+1.2} -2.0 3875.6±0.7 ^{+1.4} -1.5		D ⁰ D ⁰ π ⁰	B decays
Z(3930)	Belle	3929±5±2 + i(29±10±2)	2++	D ⁰ D ⁰ , D ⁺ D ⁻	ŶŶ
Y(3940)	Belle BaBar	3943±11±13 + i(87±22±26) 3914.3 ^{+3.8} - _{3.4} ±1.6+ i(33 ⁺¹² - ₈ ±0.60)	J++	ωJ/ψ	B decays
X(3940)	Belle	3942 ⁺⁷ -6±6 + i(37 ⁺²⁶ -15±8)	J ^{₽+}	DD*	e⁺e⁻ (recoil against J/ψ)
Y(4008)	Belle	4008±40 ⁺⁷² -28 + i(226±44 ⁺⁸⁷ -79)	1	π⁺π⁻ፓ/ψ	e⁺e⁻ (ISR)
X(4160)	Belle	4156 ⁺²⁵ -20±15+ i(139 ⁺¹¹¹ -61±21)	J [₽]	D*D*	e⁺e⁻ (recoil against J/ψ)
Y(4260)	BaBar Cleo Belle	$4259\pm8^{+8}_{-6} + i(88\pm23^{+6}_{-4})$ $4284^{+17}_{-16}\pm4 + i(73^{+39}_{-25}\pm5)$ $4247\pm12^{+17}_{-32} + i(108\pm19\pm10)$	1	π⁺π⁻Ϳ/ψ, π ^ο π ^ο Ϳ/ψ, Κ⁺Κ⁻Ϳ/ψ	e⁺e⁻ (ISR), e⁺e⁻
Y(4350)	BaBar Belle	4324±24 + i(172±33) 4361±9±9 + i(74±15±10)	1	π⁺π⁻ψ(2S)	e⁺e⁻ (ISR)
Z+(4430)	Belle	4433±4±1+ i(44 ⁺¹⁷ -13 ⁺³⁰ -11)	J٩	π ⁺ ψ(2S)	B decays
Y(4620)	Belle	4664±11±5 + i(48±15±3)	1	π⁺π⁻ψ(2S)	e⁺e⁻ (ISR)

- Much interest lately in new states observed in charmonium region: X(3872), X(3940), Y(3940), Y(4260), and Z(3930)
- X(3872) of particular interest b/c may be the first hadron-antihadron ($D^0 \overline{D}^{*0}$ + c.c.) molecule

- Much interest lately in new states observed in charmonium region: X(3872), X(3940), Y(3940), Y(4260), and Z(3930)
- X(3872) of particular interest b/c may be the first hadron-antihadron ($D^0 \overline{D}^{*0}$ + c.c.) molecule
 - need very precise mass measurement to confirm or refute
 - $\Rightarrow \overline{p}p \rightarrow X(3872)$ formation *ideal* for this

Also,...

- Study other X, Y, Z states
- Worthwhile measurements that E835 could have made but didn't...

(lack of beam time for precision scans when one didn't know exactly where to look)

- h_c mass & width, χ_c radiative-decay angular distributions, η_c' full and radiative widths,...
- ...improved limits on p
 lifetime and branching ratios (APEX),...



Charm!

PHYSICAL REVIEW D 77, 034019 (2008)

Estimate of the partial width for X(3872) into $p\bar{p}$

Eric Braaten

Physics Department, Ohio State University, Columbus, Ohio 43210, USA (Received 13 November 2007; published 25 February 2008)

We present an estimate of the partial width of X(3872) into $p\bar{p}$ under the assumption that it is a weakly bound hadronic molecule whose constituents are a superposition of the charm mesons $D^{*0}\bar{D}^0$ and $D^0\bar{D}^{*0}$. The $p\bar{p}$ partial width of X is therefore related to the cross section for $p\bar{p} \rightarrow D^{*0}\bar{D}^0$ near the threshold. That cross section at an energy well above the threshold is estimated by scaling the measured cross section for $p\bar{p} \rightarrow K^{*-}K^+$. It is extrapolated to the $D^{*0}\bar{D}^0$ threshold by taking into account the threshold resonance in the 1⁺⁺ channel. The resulting prediction for the $p\bar{p}$ partial width of X(3872) is proportional to the square root of its binding energy. For the current central value of the binding energy, the estimated partial width into $p\bar{p}$ is comparable to that of the P-wave charmonium state χ_{c1} .

 Braaten estimate of pp X(3872) coupling assuming D*D molecule

Charm!

PHYSICAL REVIEW D 77, 034019 (2008)

Estimate of the partial width for X(3872) into $p\bar{p}$

Eric Braaten

Physics Department, Ohio State University, Columbus, Ohio 43210, USA (Received 13 November 2007; published 25 February 2008)

We present an estimate of the partial width of X(3872) into $p\bar{p}$ under the assumption that it is a weakly bound hadronic molecule whose constituents are a superposition of the charm mesons $D^{*0}\bar{D}^0$ and $D^0\bar{D}^{*0}$. The $p\bar{p}$ partial width of X is therefore related to the cross section for $p\bar{p} \rightarrow D^{*0}\bar{D}^0$ near the threshold. That cross section at an energy well above the threshold is estimated by scaling the measured cross section for $p\bar{p} \rightarrow K^{*-}K^+$. It is extrapolated to the $D^{*0}\bar{D}^0$ threshold by taking into account the threshold resonance in the 1⁺⁺ channel. The resulting prediction for the $p\bar{p}$ partial width of X(3872) is proportional to the square root of its binding energy. For the current central value of the binding energy, the estimated partial width into $p\bar{p}$ is comparable to that of the P-wave charmonium state χ_{c1} .

 Braaten estimate of pp X(3872) coupling assuming D*D molecule

extrapolates from
 K*K data

Charm!

PHYSICAL REVIEW D 77, 034019 (2008)

Estimate of the partial width for X(3872) into $p\bar{p}$

Eric Braaten

Physics Department, Ohio State University, Columbus, Ohio 43210, USA (Received 13 November 2007; published 25 February 2008)

We present an estimate of the partial width of X(3872) into $p\bar{p}$ under the assumption that it is a weakly bound hadronic molecule whose constituents are a superposition of the charm mesons $D^{*0}\bar{D}^0$ and $D^0\bar{D}^{*0}$. The $p\bar{p}$ partial width of X is therefore related to the cross section for $p\bar{p} \rightarrow D^{*0}\bar{D}^0$ near the threshold. That cross section at an energy well above the threshold is estimated by scaling the measured cross section for $p\bar{p} \rightarrow K^{*-}K^+$. It is extrapolated to the $D^{*0}\bar{D}^0$ threshold by taking into account the threshold resonance in the 1⁺⁺ channel. The resulting prediction for the $p\bar{p}$ partial width of X(3872) is proportional to the square root of its binding energy. For the current central value of the binding energy, the estimated partial width into $p\bar{p}$ is comparable to that of the P-wave charmonium state χ_{c1} .

- Braaten estimate of pp X(3872) coupling assuming D*D molecule
 - extrapolates from
 K*K data
- By-product is $D^*\overline{D}$ cross section


























...and **now** for something completely different!

D. M. Kaplan, IIT

New Experiments with Antiprotons

ICL HEP Seminar 31

- Long quest at LEAR, then AD (ATRAP, ATHENA, ALPHA), to study antihydrogen and test CPT
 - e.g., is Lamb shift identical for H and \overline{H} ?
- Struggling with difficulty of combining antiprotons with positrons in a Penning trap and winding up in (or near) ground state

Antihydrogen But over 10 years ago, LEAR PS210 & FNAL E835 produced oodles of H!

Antihydrogen But over 10 years ago, LEAR PS210 & FNAL E835 produced oodles of H!

Production of antihydrogen

G. Baur^a, G. Boero^b, S. Brauksiepe^a, A. Buzzo^b, W. Eyrich^c, R. Geyer^a, D. Grzonka^a, J. Hauffe^c, K. Kilian^a, M. LoVetere^b, M. Macri^b, M. Moosburger^c, R. Nellen^a, W. Oelert^a, S. Passaggio^b, A. Pozzo^b, K. Röhrich^a, K. Sachs^a, G. Schepers^e, T. Sefzick^a, R.S. Simon^d, R. Stratmann^d, F. Stinzing^c, M. Wolke^a

^a IKP, Forschungszentrum Jülich GmbH, Germany ^b Genoa University and INFN. Italy ^c PI. Universität Erlangen-Nürnberg, Germanv ^d GSI Darmstadt. Germany ^e IKP. Universität Münster, Germanv

Received 8 December 1995; revised manuscript received 21 December 1995 Editor: L. Montanet

Abstract

Results are presented for a measurement for the production of the antihydrogen atom $\overline{H}^0 \equiv \overline{p}e^+$, the simplest atomic bound state of antimatter.

A method has been used by the PS210 collaboration at LEAR which assumes that the production of \overline{H}^0 is predominantly mediated by the e⁺e⁻-pair creation via the two-photon mechanism in the antiproton-nucleus interaction. Neutral \overline{H}^0 atoms are identified by a unique sequence of characteristics. In principle \overline{H}^0 is well suited for investigations of fundamental CPT violation studies under different forces, however, in our investigations we concentrate on the production of this antimatter object, since so far it has never been observed before.

The production of 11 antihydrogen atoms is reported including possibly 2 ± 1 background signals, the observed yield agrees with theoretical predictions. D. M. Kaplan, III



be $1.12 \pm 0.14 \pm 0.09$ pb. [S0031-9007(98)05685-3]



• Formed automatically e.g. in E835 gas-jet target, detected in "parasitic" E862





 Subsequently worked out technique to measure Lamb shift of relativistic H in flight:

• Subsequently worked out technique to measure Lamb shift of relativistic H in flight:

PHYSICAL REVIEW D

VOLUME 57, NUMBER 11

1 JUNE 1998

Measuring the antihydrogen Lamb shift with a relativistic antihydrogen beam

G. Blanford, K. Gollwitzer, M. Mandelkern, J. Schultz, G. Takei, and G. Zioulas University of California at Irvine, Irvine, California 92717

> D. C. Christian Fermilab, Batavia, Illinois 60510

> > C. T. Munger

Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309 (Received 18 December 1997; published 4 May 1998)

We propose an experiment to measure the Lamb shift and fine structure (the intervals $2s_{1/2}-2p_{1/2}$ and $2p_{1/2}-2p_{3/2}$) in antihydrogen. A sample of 10 000 antihydrogen atoms at a momentum of 8.85 GeV/*c* suffices to measure the Lamb shift to 5% and the fine structure to 1%. Atomic collisions excite antihydrogen atoms to states with n=2; field ionization in a Lorentz-transformed laboratory magnetic field then prepares a particular n=2 state, and is used again to analyze that state after it is allowed to oscillate in a region of zero field. This experiment is feasible at Fermilab. [S0556-2821(98)04711-0]

D. M. Kaplan, IIT

- Further parasitic running appears feasible
- Hope to install high-Z foil operable in Antiproton Accumulator beam halo at upcoming shutdown
- Can then assemble Lamb-shift apparatus (magnets, laser, detectors) and begin shakedown and operation

• From D. Christian:

CPT test using relativistic antihydrogen

- Antihydrogen is produced in the gas-jet target exits the Accumulator in the ground state.
 - 99 antihydrogen atoms were observed by E862 with 0 background.
- The atoms enter a 7kG magnet and a large fraction are excited to N=2 longlived Stark state by laser light.
- Atoms exit magnet & pass through a field-free region, then enter a second magnet with field 6-8 kG. The mixture of N=2 Stark states in the second magnet depends on the time spent in the field-free region, the fine structure, and the Lamb shift.
- Distribution of field ionization in the second magnet reflects probability of being in each of the three N=2 Stark states.
- Monte Carlo —> an experiment in which 100 atoms exit the first magnet in N=2,L will yield a 1% measurement of the fine structure and a 5% measurement of the Lamb shift. Assuming that only the 2S level is shifted by a CPT violating force, the 1σ sensitivity is 50 parts per billion of the 2S binding energy.

• Experimentally, unknown whether antimatter falls up or down!

• Experimentally, unknown whether antimatter falls up or down! Or whether $g - \overline{g} = 0$ or ε

- Experimentally, unknown whether antimatter falls up or down! Or whether $g \overline{g} = 0$ or ε
 - in principle a simple interferometric measurement with slow H beam [T. Phillips, Hyp. Int. 109 (1997) 357]:

- Experimentally, unknown whether antimatter falls up or down! Or whether $g \overline{g} = 0$ or ε
 - in principle a simple interferometric measurement with slow H beam [T. Phillips, Hyp. Int. 109 (1997) 357]:



- Experimentally, unknown whether antimatter falls up or down! Or whether $g \overline{g} = 0$ or ε
 - in principle a simple interferometric measurement with slow H beam [T. Phillips, Hyp. Int. 109 (1997) 357]:



- Experimentally, unknown whether antimatter falls up or down! Or whether $g \overline{g} = 0$ or ε
 - in principle a simple interferometric measurement with slow H beam [T. Phillips, Hyp. Int. 109 (1997) 357]:



 $\rightarrow \overline{g} = -g$ gives natural explanations for baryon asymmetry & dark energy

D. M. Kaplan, IIT

- Experimentally, unknown whether antimatter falls up or down! Or whether $g \overline{g} = 0$ or ε
 - in principle a simple interferometric measurement with slow H beam [T. Phillips, Hyp. Int. 109 (1997) 357]:


Antimatter Gravity

- Experimentally, unknown whether antimatter falls up or down! Or whether $g \overline{g} = 0$ or ε
 - in principle a simple interferometric measurement with slow H beam [T. Phillips, Hyp. Int. 109 (1997) 357]:



Antimatter Gravity

- Lol presented to FNAL PAC in March
- Emphasized practicality of $I^{\underline{st}} \overline{g}$ meas't to $I0^{-2}$
 - req's just 1% of 1 day's \overline{p} production
- PAC & PO (June):
 - I. interesting physics!
 - 2. but 10⁻² meas't not worthwhile (nucl. B.E.)
 - 3. need matter demo
- We're now developing techs. for 10⁻⁴ meas't & assembling matter demo



Hydrogen 2S Beam at Fermilab

Cold multichannel nozzle Excited to 2S with pulsed cathode Detected by quenching 2S, observing Lyman- α photon







D. M. Kaplan, IIT

New Experiments with Antiprotons

ICL HEP Seminar 39

Antimatter Gravity

- Requires development of deceleration techniques from 8 GeV to <20 keV:
 - MI from 8 GeV to \leq 400 MeV (TBD)
 - from ~400 MeV to 20 keV, application of µ-coolinginspired technique looks highly promising!
 - efficiency $\gtrsim 10^{-5}$ looks feasible $\Rightarrow 10^{-4} \overline{g}$ meas't in ~3 months' dedicated running
- Requires completion of antiproton deceleration/ extraction facility planned for Hbar Technologies



• From G. Jackson:



1275 W. Roosevelt Rd., Suite 130, West Chicago IL, 60185 www.hbartech.com

The HiPAT trap



- Designed to hold 1E12 antiprotons
- Designed to be portable
- Traditional superconducting solenoid requiring liquid helium for the superconductors and liquid nitrogen for the heat shield
- Good vacuum lifetime
- Comes with proton and Hlinacs for commissioning
- Still at NASA MSFC

2/21/08



A New Pbar Experiment

D. M. Kaplan, IIT

New Experiments with Antiprotons

ICL HEP Seminar

42

3

Is There an Interested Collaboration?

Is There an Interested Collaboration?

- I am drafting Lol and soliciting collaborators
 - so far:



ICL HEP Seminar 43





• Best experiment ever on hyperons, charm, and charmonia may soon be feasible at Fermilab

- Best experiment ever on hyperons, charm, and charmonia may soon be feasible at Fermilab
 - including world's most sensitive charm CPV study

- Best experiment ever on hyperons, charm, and charmonia may soon be feasible at Fermilab
 - including world's most sensitive charm CPV study
- Unique tests of CPT symmetry & antimatter gravity may be starting up soon

- Best experiment ever on hyperons, charm, and charmonia may soon be feasible at Fermilab
 - including world's most sensitive charm CPV study
- Unique tests of *CPT* symmetry & antimatter gravity may be starting up soon
- BUT...fighting uphill battle for approval!

- Best experiment ever on hyperons, charm, and charmonia may soon be feasible at Fermilab
 - including world's most sensitive charm CPV study
- Unique tests of *CPT* symmetry & antimatter gravity may be starting up soon
- BUT...fighting uphill battle for approval!
 - You can help! Want to join?

- Best experiment ever on hyperons, charm, and charmonia may soon be feasible at Fermilab
 - including world's most sensitive charm CPV study
- Unique tests of *CPT* symmetry & antimatter gravity may be starting up soon
- BUT...fighting uphill battle for approval!

You can help! Want to join?

And, please, help spread the word!

- Best experiment ever on hyperons, charm, and charmonia may soon be feasible at Fermilab
 - including world's most sensitive charm CPV study
- Unique tests of *CPT* symmetry & antimatter gravity may be starting up soon
- BUT...fighting uphill battle for approval!

You can help! Want to join?

And, please, help spread the word!

(See http://capp.iit.edu/hep/pbar)

Some HyperCP Publications:

- L. C. Lu *et al.*, "Measurement of the asymmetry in the decay $\overline{\Omega}^+ \to \overline{\Lambda} K^+ \to \overline{p} \pi^+ K^+$," Phys. Rev. Lett. **96**, 242001 (2006).
- D. Rajaram *et al.*, "Search for the Lepton-Number-Violating Decay $\Xi^- \rightarrow p\mu^-\mu^-$," Phys. Rev. Lett. **94**, 181801 (2005).
- C. G. White *et al.*, "Search for Delta $\Delta S = 2$ Nonleptonic Hyperon Decays," Phys. Rev. Lett. **94**, 101804 (2005).
- H. K. Park *et al.*, "Evidence for the Decay $\Sigma^+ \rightarrow p\mu^+\mu^-$," Phys. Rev. Lett. **94**, 021801 (2005).
- M. Huang *et al.*, "New Measurement of $\Xi^- \rightarrow \Lambda \pi^-$ Decay Parameters," Phys. Rev. Lett. **93**, 011802 (2004);
- M. J. Longo *et al.*, "High-Statistics Search for the $\Theta^+(1.54)$ Pentaquark," Phys. Rev. D 70, 111101(R) (2004);
- T. Holmstrom *et al.*, "Search for *CP* Violation in Charged-Ξ and Λ Hyperon Decays," Phys. Rev. Lett. **93**, 262001 (2005);
- Y. C. Chen *et al.*, "Measurement of the Alpha Asymmetry Parameter for the $\Omega^- \to \Lambda K^-$ Decay," Phys. Rev. D **71**, 051102(R) (2005);
- L. C. Lu *et al.*, "Observation of Parity Violation in the $\Omega^- \to \Lambda K^-$ Decay," Phys. Lett. B **617**, 11 (2005).
- R. A. Burnstein *et al.*, "HyperCP: A High-Rate Spectrometer for the Study of Charged Hyperon and Kaon Decays," Nucl. Instrum. Methods A **541**, 516 (2005).

Backup

Table 5: Summary of predicted hyperon *CP* asymmetries.

Asymm.	Mode	SM	NP	Ref.
A_{Λ}	$\Lambda o p\pi$	$\stackrel{<}{_\sim} 10^{-5}$	$\stackrel{<}{_\sim} 6 \times 10^{-4}$	[68]
$A_{\Xi\Lambda}$	$\Xi^{\mp} \to \Lambda \pi, \Lambda \to p \pi$	$\stackrel{<}{_\sim} 0.5 \times 10^{-4}$	$\leq 1.9 \times 10^{-3}$	[69]
$A_{\Omega\Lambda}$	$\Omega \to \Lambda K, \Lambda \to p\pi$	$\leq 4 \times 10^{-5}$	$\leq 8 \times 10^{-3}$	[36]
$\Delta_{\Xi\pi}$	$\Omega \to \Xi^0 \pi$	2×10^{-5}	$\leq 2\times 10^{-4}{}^*$	[35]
$\Delta_{\Lambda K}$	$\Omega \to \Lambda K$	$\leq 1 \times 10^{-5}$	$\leq 1 \times 10^{-3}$	[36]

*Once they are taken into account, large final-state interactions may increase this prediction

Backup

• Some Hyperon CP references:

- [32] A. Pais, Phys. Rev. Lett. 3, 242 (1959); O. E. Overseth and S. Pakvasa, Phys. Rev. 184, 1663 (1969); J. F. Donoghue and S. Pakvasa, Phys. Rev. Lett. 55, 162 (1985).
- [33] J. F. Donoghue, X.-G. He, S. Pakvasa, Phys. Rev. D 34, 833 (1986); X.-G. He, H. Steger, G. Valencia, Phys. Lett. B 272, 411 (1991).
- [34] G. Valencia, Proc. $\overline{p}2000$ Workshop, D. M. Kaplan and H. A. Rubin, eds., Illinois Institute of Technology, Chicago, IL 60616, USA, Aug. 3–5, 2000.
- [35] J. Tandean, G. Valencia, Phys. Lett. B 451, 382 (1999).
- [36] J. Tandean, Phys. Rev. **70**, 076005 (2004).
- [68] D. Chang, X.-G. He, and S. Pakvasa, Phys. Rev. Lett. 74, 3927 (1995).
- [69] X.-G. He, H. Murayama, S. Pakvasa, G. Valencia, Phys. Rev. D 618, 071701(R) (2000).