






Search for exotic physics with long-lived particles at ATLAS



Cristiano Sebastiani
7 June 2023, Imperial College



Large Hadron Collider



ATLAS@LHC




Standard Model

All about discovering new physics:

- ATLAS extensive measurements programme can still be considered as searches
- Almost all collider measurements to date, across 14 orders of magnitude, agree with its predictions
- ... no new physics so far...

Standard Model Production Cross Section Measurements

Status: February 2022



DM signatures

Signature based searches:

- Many expected SUSY and WIMP particles to follow shortly after the Higgs, but now increasingly disfavoured
- No hint for new discoveries to be made at the energy scales accessible to the LHC


Are there underlying assumptions in our research programme which are preventing a discovery?

it's time to explore new ideas!

ATLAS Heavy Particle Searches* - 95% CL Upper Exclusion Limits

Status: July 2022

ATLAS Preliminary
 $\int \mathcal{L} dt = (3.6 - 139) \text{ fb}^{-1}$
 $\sqrt{s} = 8, 13 \text{ TeV}$




*Only a selection of the available mass limits on new states or phenomena is shown.

†Small-radius (large-radius) jets are denoted by the letter j (J).

A new perspective

New Physics can be decoupled from electroweak scale in Dark Sector models, requiring additional low-mass mediators to explain the observed relic density with light DM (sub-GeV)



Dark Sector portals


Light mediators, HNL and ALPs must be SM singlets: options limited by SM gauge invariance

Pseudoscalar portal: $\delta_\mu \bar{\psi} \gamma^\mu \gamma^5 \psi$, $\frac{a}{f_a} F'_{\mu\nu} \tilde{F}^{\mu\nu}$ axions/ALPs

Vector portal: $\epsilon F^{\mu\nu} F'_{\mu\nu}$ ‘dark’ vector boson (A' , γ_d , Z_d) which mixes with SM photon

Higgs portal: $\kappa H^2 S^2 + \mu H^2 S$ ‘dark’ scalar boson (S) —> exotic Higgs decays

Neutrino portal: $\kappa(HL)N$ no more sterile neutrino



Feebly interacting particles are well motivated but their mass scale is unknown and are very difficult to probe at particle colliders, often lead to unconventional signatures!



Dark Sector portals

Light mediators, HNL and ALPs must be SM singlets: options limited by SM gauge invariance

Pseudoscalar portal: $\delta_u \bar{\psi} \gamma^\mu \gamma^5 \psi$ Today's study case
Vector portal: $\epsilon F^{\mu\nu} F'_{\mu\nu}$ 'dark' vector boson (A' , γ_d , Z_d) which mixes with SM photon

Higgs portal: $\kappa H^2 S^2 + \mu H^2 S$ 'dark' scalar boson (S) \rightarrow exotic Higgs decays


Neutrino portal: $\bar{\nu}_L H \Delta M$ no more sterile neutrino
New particles can be long-lived:
observed lifetime is governed by an
exponential defined by the proper lifetime $c\tau$



Exotic among exotics?

Search for new BSM physics at LHC with exotic signatures:

- Standard decays
- **Unconventional signatures**: long time-of-flight, anomalous energy deposits, displaced secondary vertices...
- Detector-stable particles



ATLAS and CMS are not optimised for this kind of signals

Unusual and unique signatures are extremely challenging to probe:

TRIGGER

Anomalous signatures not associated with standard activity in the detector require the development of dedicated triggers!

RECONSTRUCTION

Object identification and reconstruction algorithms are to be updated to include non-standard tracks and energy deposits

NON-COLLISION BACKGROUND


Unconventional signatures have unconventional backgrounds, from detector noise to non-collision physics events

Unconventional backgrounds

Develop new techniques and ideas to reduce very unconventional background sources


Cosmic muons

- Muon bundles from cosmic-ray that cross the detector in-time with the pp collision event




Beam induced background (BIB)

- Machine/beam-induced backgrounds from upstream proton losses in the LHC, from inelastic collisions with residual gas, beam halo cleaning losses or beam-gas scattering



BIB event




Run: 300487

Event: 445566353

2016-05-30 00:29:53 CEST

EXOT-2019-05

Unconventional signatures



LLPs could lead to a plethora of new signatures. Experimental approach depending on...

- Charge and decay products
- Interaction with specific sub-detector
- Displacement of the decay

LLP guidebook

- Find a new signature yet unexplored in ATLAS broad LLP programme
- Interaction with the LHC community and theorists to exchange ideas
- Use the detector at its best: bespoke strategies for triggers, reconstruction and calibration objects
- Exploit new state of the art techniques, like Machine Learning tools
- Enjoy the journey :)



MY WORK-LIFE BALANCE



NEW LLP ANALYSES


A long-lived dark photon search

Exploit ‘large’ Higgs cross-section to probe events with very small epsilon values → Long-lived particles (LLP)



Current constraints


Constraints on visible dark photon decays are expressed in the parameter space epsilon ϵ versus dark photon mass $m(A')$



Region explorable at particle
colliders via Higgs portal

Search for long-lived dark photons

Search for light long-lived neutral particles decaying into collimated jet structures of leptons or light hadrons



Muonic decay


Collimated bundle of muon without track in the inner detector (no jets)

**Difficult to trigger: low-pt muon
Cosmic-ray muons background**

$$\Delta\Phi \simeq \frac{m_{\gamma d}}{E_{\gamma d}} << 1$$


Outgoing products
may be collimated

Vector portal



For small couplings
very displaced decays

$$c\tau = \frac{1}{\Gamma_{tot}} \propto \frac{1}{\epsilon^2}$$




Calorimeter decay

**Displaced jet with most of energy
deposit in the HCAL (no muons)**

**Very high background from rare QCD
events and few handles to play with**




Dark-photon candidate



Run: 303266
Event: 1584619053
2016-07-04 04:57:58 CEST
EXOT-2019-05

Trigger for muonic dark-photon

Dedicated triggers are used to select events with highly collimated muons with low activity in the ID



Analysis strategy

- Dedicated triggers: NarrowScan (muonic) and CaloRatio (hadronic)
- Two production modes: ggH and WH
 - Six channels: muonic-muonic (ggH), muonic-calorimeter (ggH+WH), one-calorimeter exclusive (WH) and calorimeter-calorimeter (ggH+WH)
- Backgrounds: rare QCD events, cosmic-ray muons and beam induced background (BIB)
 - Advanced NN-based taggers according to DPJ type
- Data-driven ABCD method to estimate dominant QCD background in the signal regions

Event Selection

1. Trigger
2. Quality Cuts



DPJ Reconstruction

1. Muonic-Muonic
2. Muonic-Calorimeter
3. One-Calorimeter
4. Calorimeter-Calorimeter




Taggers Cuts



ABCD method


Benchmark models

DPJ



Hidden Abelian Higgs Model
[HAHM] arXiv:1312.4992


DPJ





Falkovsky, Ruderman, Volansky,
Zupan [FRVZ] arXiv:1002.2952

A low-level input NN

Extract low level information from the ATLAS calorimeter from a single jet in either 3D images or graphs



Calorimeter sampling is sensitive to the dark photon displacement






Tool to reject QCD main background, will allow an exclusion on hadronic-LJ channels!

A closer look on NN methods



The ATLAS detector orthogonal view



Let's exploit the calorimeter granularity to parametrise the energy deposits: x, y, z, energy

3D jet images:

- Train a CNN
- Very sparse images -> sub-optimal

Graphs:

- Train a fully optimised GNN
- Small cloud space objects
- Efficient and easy to manipulate


Additional higher level variable can be added as features to further improve the network performance, although the goal is to have them already 'learned' by the network by using only the low level inputs

Project developed within the **MUCCA** consortium - Multi-disciplinary Use Cases for Convergent new Approaches to AI explainability, with M. D'Onofrio and J. Carmignani —> for more info: [CHIST-ERA-video](#)

Performances



The GNN model out-performed the CNN model on all **performance metrics** tested at same signal efficiency score




As expected graph dataset are proven very effective for classification of sparse image of HEP calorimeter detectors!
GNN will be used in Run3 searches!

Dark-photon explainability



- ▶ How does a signal look like?




- ▶ How does the NN understands true positive (signal), true negative or false positive ?

▶ Work in progress!

(True Negative)

➤ Sig 1 Bkg 0

True label: 0, predicted label: 0, predicted prob: 0.01




Input Graph

Trac-In

(False Positive)

True label: 0, predicted label: 1, predicted prob: 0.76




Input Graph


Trac-In

- ❖ Proponents for Bkg while opponents for Sig which is sensibly expected


[Proponent] Label:0



[Proponent] Label:0




[Proponent] Label:0




[Proponent] Label:0


[Opponent] Label:1



[Opponent] Label:1




[Opponent] Label:1




[Opponent] Label:1


[Opponent] Label:1





[Opponent] Label:1



[Opponent] Label:1





[Opponent] Label:1



Background estimation

Data-driven estimation of residual multi-jet background events in signal region defined by a pair of cuts of two uncorrelated variables for the background


- Single source of background (remove all BIB and Cosmics, leave only rare QCD events)
- No correlation between variables
- Small signal contamination in CRs



$$N_A^{exp} = \frac{N_D \cdot N_B}{N_C} \text{Var 1}$$

Results


No excess found! Exclusion limits are presented as exclusion regions in the $(\varepsilon, \text{myd})$ plane in the context of the Vector portal model as a function of BR ($h \rightarrow$ dark sector)




Combination of ggH and WH hadronic channels to improve results!

First time exclusion in the fully electron channel


Muonic exclusion of $\text{Br}(h \rightarrow \text{dark sector})$ down to 0.1%!
 $\rightarrow \text{Br}(h \rightarrow 4\nu) \sim 0.1\%$!



Significant analysis improvements (CNN taggers) and new topologies (WH) allowed for an exclusion of hadronic decays




First use in ATLAS searches of a Convolutional Neural Network based tagger trained on low-level inputs (3D jet images from calorimetric clusters)




Results


No excess found! Exclusion limits are presented as exclusion regions in the $(\varepsilon, \text{myd})$ plane in the context of the Vector portal model as a function of BR ($h \rightarrow$ dark sector)




Combination of ggH and WH hadronic channels to improve results!




Muonic exclusion of $\text{Br}(h \rightarrow \text{dark sector})$ down to 0.1%!
 $\rightarrow \text{Br}(h \rightarrow 4\nu) \sim 0.1\%$!



First time exclusion of HAHM model with dark-photon jet signature





First use in ATLAS searches of a Convolutional Neural Network based tagger trained on low-level inputs (3D jet images from calorimetric clusters)



mono-jet reinterpretation

Both dark photons can decay outside ATLAS...


[ATL-PHYS-PUB-2021-](#)



Monojet is becoming a precision measurement with a huge interpretation potential:

- Model-independent limits on visible cross-section
- Higgs to invisible BR upper limit (95%CL obs. 0.50)
- Countless possible NP interpretations

Where to go now?



Full Run2 prompt analysis effort restarted, results for LHCb:

- Extend $m_{\gamma d}$ mass range to $[0.01;10] \text{ GeV}$

Displaced analysis to be improved with VBF signature:

- Significantly improve ele results ($m_{\gamma d} < 280 \text{ MeV}$)

Analyses are designed to aim for a final combination of prompt + displaced (ggF+WH+VBF)! Run2 legacy result of FRVZ and HAHM models

Where to go in Run3:


- Go TLA+PEB for muonic signature
- Additional Higgs production mechanism for prompt analysis
- Explore mono-LJ (LJ + MET) signature to fill the gaps between prompt and combined

Not only dark-photons in ATLAS...

Snapshot of the most recent ATLAS LLP results..


ATLAS Long-lived Particle Searches* - 95% CL Exclusion

Status: July 2022



*Only a selection of the available lifetime limits is shown.

Analyses reach into the lifetime space




From the past year...

- (Mar 23) CONF-2023-018 Micro displaced muons
- (Jul 23) CONF-2022-051 Displaced Higgs in electron or photon
- (Jan 23) SUSY-2018-13 DV+Jets
- (Jul 22) SUSY-2019-14 Non-pointing and delayed photons
- (Jun 22) SUSY-2018-19 Disappearing track
- (May 22) EXOT-2019-05 Displaced lepton-jets
- (May 22) EXOT-2018-54 Multi-charged particles
- (Apr 22) SUSY-2018-42 Pixel dE/dx

Dark Sector portals


Light mediators, HNL and ALPs must be SM singlets: options limited by SM gauge invariance

Pseudoscalar portal: $\delta_\mu \bar{\psi} \gamma^\mu \gamma^5 \psi$, $\frac{a}{f_a} F'_{\mu\nu} \tilde{F}^{\mu\nu}$ **axions/ALPs**



Are we covering the whole phase space accessible at the LHC?






Decay rates for a NP scale at 1 TeV and effective coefficients set to 1



Nice complementarity between LHC, beam-dump and astrophysics boundaries. Together they can explore a large fraction of the axion regions


Prompt axion searches

Many searches looking for prompt ALPs in a great effort to systematically cover all production and decay channels!



Displaced axion searches


Still plenty of unexplored displaced ALP scenarios, many possibilities for synergies and reinterpretations



It's time to develop new ideas and explore all blind spot left in the ATLAS LLP programme!

New idea: a search for long-lived axions

- Axion particle produced in association with a Z boson from Higgs decay
- Signature with two leptons and two collimated photons (one if signal photons are collimated enough)
- Main backgrounds from Z+jet and Zy



2 close-by photons


preferred parameter space where mu g-2 anomaly can be explained

axion decay rate:

$$\Gamma(a \rightarrow \gamma\gamma) \equiv \frac{4\pi\alpha^2 m_a^3}{\Lambda^2} |C_{\gamma\gamma}^{\text{eff}}|^2$$

axion displaced region
From few mm to stable




[arXiv:1708.00443](https://arxiv.org/abs/1708.00443)




ALPs pseudo-scalar particles: mass range 0.1-35 GeV.

Current searches

Non-pointing photon search [SUSY-2019-14](#) with ID di-photon vertexing [SUSY-2020-28](#)





- Analysis optimised for **high-mass** LLP mediator that decays before **ECAL** with a MET signature
- New search plans to target **low-mass** LLP mediator that decays within **ECAL** or **HCAL**



Probing displaced photons


Target key discriminants for long-lived photons: anomalous shower shape, delayed timing and secondary vertices.

Calo images and graph to reconstruct decays within calorimeter from shower shapes




Calo vertexing to reconstruct secondary vertex within the ID

Photon timing to target non-pointing photons



Also more complete models

Long-lived particles are present in many BSM theories like SUSY and heavy neutrinos




Specific theories can suggest new signatures to explore. LLPs are well-motivated in many BSM models (due to weak couplings or small mass splitting).

Results are presented for representative benchmark scenarios, giving the possibility to reinterpret in a different model.


Unconventional SUSY models

Disappearing tracks: Search for long-lived particle decaying within the inner tracker



- MET trigger to select events with jets and ‘disappearing tracks’ (due to suppressed interaction or low-pT)
- ‘Disappearing track’: 4-hit pixel tracks with no hits in the silicon tracker (SCT) and < 5 GeV of energy deposits in calo $dR < 0.2$
- Rare SM backgrounds from charged lepton scattering and combinatorial fakes



Targets very compressed SUSY scenarios (...and various DM models)




Electroweak (EWK)
production




Strong production

Disappearing tracks

Exclusion limits at 95% CL for the EWK channel fo the AMSB model, neutralino mass vs lifetime



Exclusion for 660 GeV wino, best limit for wino-like neutralino (460 GeV in early Run2)



Exclusion for 210 GeV Higgsino, best limit for higgsino-like neutralino (155 GeV in early Run2)

Outlook

- Long-lived particle searches have been so far discontinuous in ATLAS: from the portal framework we can identify benchmark models for a systematic investigation of the hidden sectors
- A snapshot of the most recent results from ATLAS have been shown, but many more analyses in progress using the full Run-2 dataset
- Run3 with a larger dataset and new exciting improvements will enhance the discovery potential for Long-lived signatures
- Many unexplored synergies and overlaps across searches that could be exploited
 - Harmonise common models where possible or promote multiple interpretations
 - Encourage combinations and summary plots to spot uncovered regions

